

Tutorial session 2: Bacterial cell structure


Learning objectives:

- Identify bacterial shapes and arrangements from microscopic observations.
- Compare the cell wall structure of Gram-positive and Gram-negative bacteria.
- Explain the role of the bacterial cell wall in maintaining cell integrity under osmotic stress.

Exercise 1: Bacterial shape and arrangement

Bacteria observed under the light microscope after simple staining may show different shapes and cellular arrangements. These characteristics are important for bacterial identification.

Task: Examine the image below and indicate the shape and arrangement of each bacterium.



- | | |
|--------|---------|
| 1..... | 2 |
| 3..... | 4..... |
| 5..... | 6..... |
| 7..... | 8..... |
| 9..... | 10..... |

Exercise 2: Study of the bacterial cell wall

The bacterial cell wall is a fundamental structure that differs significantly between Gram-positive and Gram-negative bacteria.

Two bacteria are studied:

- Bacterium A: Gram-positive
- Bacterium B: Gram-negative

Questions:

1. Compare the thickness of the cell wall in Gram-positive and Gram-negative bacteria.

.....

2. List the main chemical components of the bacterial cell wall.

.....
.....
.....
.....

3. Indicate the elements specific to:

- Gram-positive bacteria

.....
.....

- Gram-negative bacteria

.....
.....

4. Explain why Gram staining allows differentiation between these two groups.

.....
.....
.....

Exercise 3: The role of the bacterial cell wall

Two bacterial species are studied **separately**:

- **Bacterium 1:** *Bacillus subtilis* (Gram-positive)
- **Bacterium 2:** *Escherichia coli* (Gram-negative)

Each bacterium is suspended in different media, under controlled conditions.

Gram-positive bacterium (*Bacillus subtilis*)

The same bacterial suspension is divided into three separate test tubes.

Tube	Medium	Additional treatment
1	Distilled water (hypotonic)	None
2	Isotonic solution (physiological solution / sucrose)	Lysozyme
3	Distilled water (hypotonic)	Lysozyme

Note: Target of lysozyme : Lysozyme degrades β (1–4) glycosidic bonds between NAM and NAG in peptidoglycan

Observed results

- Tube 1: Cells keep their normal rod shape.
- Tube 2: Cells become spherical.
- Tube 3: Cells swell and burst.

Questions

1. Why do the bacteria not lyse in tube 1 despite the hypotonic medium?

.....

.....

.....

2. Why do the bacteria become spherical in tube 2?

.....

.....

.....

3. Why does lysis occur only in tube 3?

.....

.....

.....

4. What conclusion can be drawn about the role of the cell wall?

.....

.....

.....

Gram-negative bacterium (*Escherichia coli*)

The experiment is repeated using *E. coli*.

Tube	Medium	Additional treatment
4	Isotonic solution	Lysozyme
5	Isotonic solution	Lysozyme + EDTA

EDTA (ethylenediaminetetraacetic acid) is an agent that destabilizes the outer membrane of Gram-negative bacteria.

Observed results

- Tube 4: No visible change in cell shape.
- Tube 5: Cells become spherical but do not burst.

Questions

1. Why does lysozyme alone have no effect in tube 4?

.....

.....

.....

2. Why are the cells spherical but still intact in tube 5??

.....

.....

.....

Synthesis and interpretation

Answer the following questions using the previous exercises:

1. Define protoplast and spheroplast.

.....

.....

2. Explain why bacteria can survive in hypotonic environments.

.....

3. Complete the following sentence:

“The bacterial cell wall, mainly composed of....., plays an essential role in maintaining and protecting the cell against”